

VARIATION OF QUASICONFORMAL MAPPINGS ON LINES

LEONID V. KOVALEV AND JANI ONNINEN

ABSTRACT. We obtain improved regularity of homeomorphic solutions of the reduced Beltrami equation, as compared to the standard Beltrami equation. Such an improvement is not possible in terms of Hölder or Sobolev regularity; instead, our results concern the generalized variation of restrictions to lines. Specifically, we prove that the restriction to any line segment has finite p -variation for all $p > 1$ but not necessarily for $p = 1$.

1. INTRODUCTION

A key property of Sobolev functions in Euclidean spaces is their absolute continuity on almost every line parallel to the coordinate axes. The restrictions to arbitrary lines need not be even bounded for functions in Sobolev spaces $W^{1,s}$, $1 \leq s \leq n$. However, for $s > n$ the restriction of a Sobolev function to *any* line has finite p -variation with $p = s/(s - n + 1)$, see Remark 2.1. Here we refer to the generalized variation [19, 23, 24], which is defined as follows.

For distinct points $a, b \in \mathbb{R}^n$ we write $[a, b] = \{(1 - t)a + tb : 0 \leq t \leq 1\}$ and call $[a, b]$ the line segment with the endpoints a and b . Any partition $0 = t_0 < t_1 < \dots < t_N = 1$ induces a partition of $[a, b]$ by the rule $a_j = a + t_j(b - a)$, $j = 0, \dots, N$.

Definition 1.1. Let $\phi : [0, \infty) \rightarrow [0, \infty)$ be a convex strictly increasing function such that $\phi(0) = 0$. A mapping f from a line segment $[a, b] \subset \mathbb{R}^n$ into $\mathbb{R}^m$ has *finite ϕ -variation* on $[a, b]$ if

$$\text{var}_{[a,b]}(f; \phi) := \sup \sum_{j=1}^N \phi(|f(a_j) - f(a_{j-1})|) < \infty,$$

Date: January 9, 2009.

2000 Mathematics Subject Classification. Primary 30C62; Secondary 26A45, 30C65, 26B30.

Key words and phrases. Bounded variation, generalized variation, quasiconformal mappings, quaternions.

Kovalev was supported by the NSF grant DMS-0700549.

Onninen was supported by the NSF grant DMS-0701059.

where the supremum is taken over all partitions $(a_j)_{j=0}^N$ of $[a, b]$ and over all $N \geq 1$.

If f is a mapping from a domain $\Omega \subset \mathbb{R}^n$ into $\mathbb{R}^m$, then we say that f has *finite ϕ -variation on lines* if its restriction to any compact line segment contained in Ω has finite ϕ -variation. When $\phi(t) = t^p$, we speak of p -variation (or simply variation if $p = 1$) and write $\text{var}_{[a,b]}(f; p)$.

We are primarily interested in the variation of quasiconformal mappings $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$, where $n \geq 2$. Recall that a sense-preserving homeomorphism $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is said to be *quasiconformal* if there exists $H < \infty$ such that

$$(1.1) \quad \limsup_{r \rightarrow 0} \frac{\max_{|x-a|=r} |f(x) - f(a)|}{\min_{|y-a|=r} |f(y) - f(a)|} \leq H, \quad \text{for every } a \in \mathbb{R}^n.$$

It is a well-known result of Gehring [9] that such mappings are absolutely continuous on almost every line (ACL). This was recently extended to Ahlfors regular metric spaces by Balogh, Koskela and Rogovin [4]. The ACL property makes it possible to give an analytic definition of quasiconformal mappings.

Definition 1.2. A homeomorphism $f \in W_{\text{loc}}^{1,n}(\mathbb{R}^n; \mathbb{R}^n)$ is K -quasiconformal, $1 \leq K < \infty$, if it satisfies the distortion inequality

$$(1.2) \quad \|Df(x)\|^n \leq KJ(x, f) \quad \text{a.e.}$$

Here $\|Df(x)\|$ stands for the norm of the differential matrix and $J(x, f)$ for the Jacobian determinant. A mapping $f \in W_{\text{loc}}^{1,n}(\mathbb{R}^n; \mathbb{R}^n)$ (not necessarily homeomorphism) satisfying (1.2) is called K -quasiregular [20, 21].

Our study of the variation of quasiconformal mappings on lines grew out of [14] where it was proved that the ordinary differential equation $\dot{x} = f(x)$ has unique local solutions outside of $f^{-1}(0)$ provided that f is quasiconformal and has bounded variation on C^1 -smooth curves. By Gehring's theorem [10] quasiconformal mappings are locally in $W^{1,s}$ for some $s > n$ and therefore have finite p -variation on lines for some $p < n$. (In fact, any homeomorphism of class $W^{1,n}$ has finite n -variation on lines [18, Thm 4.3]). In the opposite direction, a theorem of Bishop [5, Thm 1.1] implies that for any $p < n$ there is a quasiconformal mapping $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$ such that the image of some line segment under f has Hausdorff dimension greater than p . Clearly, such f has infinite p -variation on this segment.

Interestingly, some classes of quasiconformal mappings exhibit much higher regularity along lines and smooth curves than their Sobolev or Hölder regularity would suggest.

Definition 1.3. A mapping $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is called δ -monotone, $0 < \delta \leq 1$, if for every $a, b \in \mathbb{R}^n$

$$(1.3) \quad \langle f(a) - f(b), a - b \rangle \leq \delta |f(a) - f(b)| |a - b|.$$

Any nonconstant δ -monotone mapping is quasiconformal [16, Cor. 7]. For example, the radial stretch $f(x) = |x|^{\alpha-1}x$, where $\alpha > 0$, is δ -monotone for some $\delta = \delta(\alpha)$. This mapping is locally Hölder continuous with exponent $\min\{\alpha, 1\}$, which can be arbitrarily close to 0. This shows that δ -monotone mappings are no more regular on the Hölder and Sobolev scales than general quasiconformal mappings. However, they have bounded variation on C^1 -smooth curves, see [2, Thm. 3.11.7] and [14, Thm. 1.10]. In particular,

$$(1.4) \quad \text{var}_{[a,b]}(f; 1) < \infty \quad \text{if } a, b \in \mathbb{R}^n.$$

When $n = 2$, we often identify $\mathbb{R}^n$ with $\mathbb{C}$ and use the complex derivatives f_z and $f_{\bar{z}}$. Then the inequality (1.2) reads as

$$(1.5) \quad |f_{\bar{z}}| \leq k |f_z| \quad \text{a.e., where } k = \frac{K-1}{K+1}$$

A δ -monotone mapping $f: \mathbb{C} \rightarrow \mathbb{C}$ satisfies the stronger, *reduced* distortion inequality

$$(1.6) \quad |f_{\bar{z}}| \leq k \operatorname{Re} f_z \quad \text{a.e. in } \mathbb{C},$$

for some constant $0 < k < 1$ (Theorem 3.11.6 [2]). The converse is false: for instance, $f(z) = iz$ satisfies (1.6) but is not δ -monotone.

Definition 1.4. A homeomorphism $f \in W_{\text{loc}}^{1,2}(\mathbb{C}; \mathbb{C})$ is called *reduced quasiconformal* if it satisfies (1.6).

Inequality (1.6) implies that f is a solution of the *reduced Beltrami equation*

$$(1.7) \quad f_{\bar{z}} = \lambda(z) \operatorname{Re} f_z \quad \text{a.e. in } \mathbb{C}$$

where $|\lambda(z)| \leq k$. Conversely, it is shown in [2, Thm 6.3.2] that any homeomorphic solution of (1.7) has constant sign of $\operatorname{Re} f_z$. Therefore such solutions satisfy (1.6) up to a change of sign.

Unlike quasiconformality, the properties (1.3) and (1.6) are preserved under addition. Both of these classes arise naturally in the theory of elliptic partial differential equations [1, 2, 3]. Our first result shows that (1.4) cannot be extended to reduced quasiconformal mappings.

Theorem 1.5. *For every $k \in (0, 1)$ there exists a reduced quasiconformal mapping $f: \mathbb{C} \rightarrow \mathbb{C}$ that satisfies (1.6) but does not have bounded variation on any nontrivial interval $[a, b] \subset \mathbb{R}$. Furthermore, f can be chosen so that $f_{\bar{z}}, \operatorname{Re} f_z \in L^\infty(\mathbb{C})$ and $\operatorname{Re} f(x) = x$ for all $x \in \mathbb{R}$.*

In other words, f in Theorem 1.5 maps $\mathbb{R}$ into a curve that is nowhere locally rectifiable. Although such examples of (non-reduced) quasiconformal mappings were known for a long time [22], our mapping f seems to be the first one given by an explicit analytic expression, see (4.1). The additive property of reduced quasiconformal mappings allows us to derive the following result from Theorem 1.5.

Corollary 1.6. *For any countable family of parallel lines $\{L_j: j = 1, 2, \dots\}$ in $\mathbb{C}$ there exists a quasiconformal mapping $f: \mathbb{C} \rightarrow \mathbb{C}$ such that any non-trivial subarc of $f(L_j)$ is unrectifiable for every j .*

Corollary 1.6 exhibits a quasiconformal mapping with irregular behavior on a relatively large set. The authors of [6] asked (Question 4.4) whether for any set $E \subset \mathbb{C}$ of planar measure zero there is a quasiconformal mapping $f: \mathbb{C} \rightarrow \mathbb{C}$ such that the volume derivative $\lim_{r \rightarrow 0} \frac{|f(B(x,r))|}{|B(x,r)|}$ is infinite at every point of E . While the singular behaviour of f in Corollary 1.6 is of different nature, the additive property of reduced quasiconformal mappings can be potentially useful in creating mappings with a large set of infinite volume derivative.

Our second main result shows that, Theorem 1.5 notwithstanding, reduced quasiconformal mappings are much more regular on lines than general quasiconformal mappings. In particular, they have finite p -variation for any $p > 1$.

Theorem 1.7. *Let $f: \mathbb{C} \rightarrow \mathbb{C}$ is a reduced quasiconformal mapping. Then for any $q > 1$ the mapping f has finite ϕ -variation on lines with*

$$(1.8) \quad \phi(t) = \frac{t}{(\log(e + 1/t))^q} \quad \text{if } t > 0 \quad \text{and} \quad \phi(0) = 0.$$

The conclusion of Theorem 1.7 is false for $0 \leq q < 1/2$, see Remark 4.1. The gap between exponents $1/2$ and 1 remains open.

Question 1.8. What is the smallest value of q for which the conclusion of Theorem 1.7 holds?

Since δ -monotone mappings exist in any dimension $n \geq 2$, one may ask whether it is possible to extend the definition of reduced quasiconformal mappings to higher dimensions. This question is addressed in section 5, where we use quaternions to define reduced quasiconformal mappings in four dimensions, and extend Theorem 1.7 to them.

Question 1.9. Is there a natural analogue of reduced quasiconformal mappings in dimensions other than 2 and 4?

2. PRELIMINARIES

In this section we first estimate the p -variation of Sobolev functions of lines. Although this result is probably known we give a proof for the sake of completeness. Later in the section we define quasisymmetric and monotone mappings and introduce some relevant notation. In this paper Ω stands for a domain in $\mathbb{R}^n$.

Proposition 2.1. *Let $u \in W^{1,s}(\Omega)$, $s > n$. Then the restriction of u to any closed line segment $I \subset \Omega$ has finite p -variation with $p = s/(s - n + 1)$.*

The Morrey-Sobolev embedding theorem states that $W^{1,s}(\Omega) \subset C_{\text{loc}}^\alpha(\Omega)$ with $\alpha = 1 - n/s$. Clearly, any function $u \in C_{\text{loc}}^\alpha(\Omega)$ has finite p -variation on lines with $p = 1/\alpha = s/(s - n)$. However, Proposition 2.1 gives a better value of p . Its proof requires the following lemma.

Lemma 2.2. *Let I be a line segment partitioned into smaller segments I_m , $m = 1, 2, \dots, M$. For any mapping $f: I \rightarrow \mathbb{R}^n$ we have*

$$(2.1) \quad \text{var}_I(f; \phi) \leq \sum_{m=1}^M \text{var}_{I_m}(f; \phi) + (M - 1) \text{osc}_I f.$$

Proof. Fix a partition $(a_j)_{j=0}^N$ of I . We divide the set of indices as follows.

$$E = \{j = 1, \dots, N : [a_{j-1}, a_j] \subset I_m \text{ for some } m\}, \quad F = \{1, \dots, N\} \setminus E.$$

Then

$$\sum_{j \in E} \phi(|f(a_j) - f(a_{j-1})|) \leq \sum_{m=1}^M \text{var}_{I_m}(f; \phi)$$

and

$$\sum_{j \in F} \phi(|f(a_j) - f(a_{j-1})|) \leq (M - 1) \text{osc}_I f.$$

□

Proof of Proposition 2.1. Dividing I into subintervals and using Lemma 2.2 we may reduce our task to the case

$$\text{diam } I < \text{dist}(I, \partial\Omega).$$

Let $(a_j)_{j=0}^N$ be a partition of I . For $j = 1, \dots, N$ let $\overline{B}_j$ be the closed ball with segment $[a_{j-1}, a_j]$ as a diameter. Morrey's inequality [7, p. 143] yields

$$\text{osc}_{\overline{B}_j} u \leq C |a_j - a_{j-1}|^{1-n/s} \left(\int_{B_j} |\nabla u(x)|^s dx \right)^{1/s}.$$

Raising to the power p and noticing that $(1 - n/s)p = 1 - p/s$ we arrive at

$$(\text{osc}_{\overline{B}_j} u)^p \leq C |a_j - a_{j-1}|^{1-p/s} \left(\int_{B_j} |\nabla u(x)|^s dx \right)^{p/s}.$$

Summing over j and applying Hölder's inequality we obtain

$$(2.2) \quad \sum_{j=1}^N (\text{osc}_{\overline{B}_j} u)^p \leq C \left(\sum_{j=1}^N |a_j - a_{j-1}| \right)^{1-p/s} \left(\sum_{j=1}^N \int_{B_j} |\nabla u(x)|^s dx \right)^{p/s}.$$

Therefore

$$\text{var}_I(u; p) \leq C (\text{diam } I)^{1-p/s} \left(\int_{\Omega} |\nabla u(x)|^s dx \right)^{p/s}$$

as desired. □

Definition 2.3. Let $\eta: [0, \infty) \rightarrow [0, \infty)$ be a homeomorphism. An injective mapping $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is η -quasisymmetric if

$$\frac{|f(c) - f(a)|}{|f(b) - f(a)|} \leq \eta \left(\frac{|c - a|}{|b - a|} \right)$$

for any distinct points $a, b, c \in \mathbb{R}^n$. The function η is called a modulus of quasisymmetry of f .

It is well-known that a mapping $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is quasiconformal if and only if it is sense-preserving and quasisymmetric [11].

Given a mapping $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$, where $\mathbb{R}^n \subset \mathbb{R}^n$, we define the modulus of monotonicity $\Delta_f: \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ by the rule

$$(2.3) \quad \Delta_f(a, b) = \begin{cases} \left\langle f(a) - f(b), \frac{a-b}{|a-b|} \right\rangle & \text{if } a \neq b \\ 0 & \text{if } a = b \end{cases}$$

Clearly $|\Delta_f(a, b)| \leq |f(a) - f(b)|$. By definition, f is a monotone mapping if $\Delta_f(a, b) \geq 0$ for all $a, b \in \mathbb{R}^n$, and is strictly monotone if $\Delta_f(a, b) > 0$ unless $a = b$. Any reduced quasiconformal is monotone by (1.9) in [14]. Also, f is δ -monotone if and only if $\Delta_f(a, b) \geq \delta |f(a) - f(b)|$ for all $a, b \in \mathbb{R}^n$. When $n = 2$, the modulus of monotonicity can be expressed in complex notation:

$$\Delta_f(a, b) = \operatorname{Re} \left(\frac{f(a) - f(b)}{a - b} \right) |a - b|.$$

3. GENERALIZED VARIATION ON LINES: PROOF OF THEOREM 1.7

We will obtain Theorem 1.7 as a consequence of the following result.

Theorem 3.1. *Let $f: \Omega \rightarrow \mathbb{R}^n$ be a mapping and suppose that there is a homeomorphism $\eta: [0, \infty) \rightarrow [0, \infty)$ such that*

$$(3.1) \quad |f(c) - f(a)| \leq \frac{|c - a|}{|b - a|} |f(b) - f(a)| + \eta \left(\frac{|c - a|}{|b - a|} \right) \Delta_f(a, b)$$

for any distinct points $a, b, c \in \Omega$. Then for any $q > 1$ the mapping f has finite ϕ -variation on lines with ϕ as in (1.8).

Before proving Theorem 3.1 we derive Theorem 1.7 from it.

Proof of Theorem 1.7. Let $f: \mathbb{C} \rightarrow \mathbb{C}$ be a reduced quasiconformal mapping. We may assume that f is nonlinear. For any $\lambda \in \mathbb{R}$ the mapping $f^\lambda(z) = f(z) + i\lambda z$ also satisfies the reduced distortion inequality 1.6 with the same constant k as f . By [13, Cor. 1.5] f^λ is a homeomorphism. Therefore, f^λ is K -quasiconformal with K independent of λ . Since quasiconformality implies quasisymmetry in $\mathbb{C}$ [11, Thm 11.14], there is a homeomorphism $\eta: [0, \infty) \rightarrow [0, \infty)$ such that f^λ is η -quasisymmetric in $\mathbb{C}$ for

all $\lambda \in \mathbb{R}$. Given distinct points $a, b, c \in \mathbb{R}$, let $\lambda = -\operatorname{Im} \frac{f(b)-f(a)}{b-a}$, so that $|f^\lambda(b) - f^\lambda(a)| = \Delta_f(a, b)$. Since f^λ is η -quasisymmetric, we have

$$|f^\lambda(c) - f^\lambda(a)| \leq \eta \left(\frac{|c-a|}{|b-a|} \right) |f^\lambda(b) - f^\lambda(a)|,$$

from which (3.1) follows by means of the triangle inequality. It remains to apply Theorem 3.1 to f . $\square$

Our proof of Theorem 3.1 is based on two lemmas.

Lemma 3.2. *Let $f: \Omega \rightarrow \mathbb{R}^n$ be as in Theorem 3.1. Given distinct points $a, b \in \Omega$ and a partition $0 = t_0 < t_1 < \dots < t_N = 1$, let $a_j = a + t_j(b-a)$, $j = 0, \dots, N$. If $a_j \in \Omega$ for all j , then*

$$(3.2) \quad \sum_{j=1}^N |f(a_j) - f(a_{j-1})| \leq C \log(N+1) |f(a) - f(b)|$$

where the constant C depends only on η in (3.1).

Proof. It suffices to prove (3.2) for $N = 2^m$. For $j = 1, \dots, 2^{m-1}$ we apply (3.1) to the points $a_{2j}, a_{2j-1}, a_{2j-2}$ and find that

$$|f(a_{2j}) - f(a_{2j-1})| \leq \frac{|a_{2j} - a_{2j-1}|}{|a_{2j} - a_{2j-2}|} |f(a_{2j}) - f(a_{2j-2})| + \eta(1) \Delta_f(a_{2j}, a_{2j-2})$$

and

$$|f(a_{2j-1}) - f(a_{2j-2})| \leq \frac{|a_{2j-1} - a_{2j-2}|}{|a_{2j} - a_{2j-2}|} |f(a_{2j}) - f(a_{2j-2})| + \eta(1) \Delta_f(a_{2j}, a_{2j-2}).$$

Adding these inequalities we obtain

$$\begin{aligned} |f(a_{2j}) - f(a_{2j-1})| + |f(a_{2j-1}) - f(a_{2j-2})| \\ \leq |f(a_{2j}) - f(a_{2j-2})| + 2\eta(1) \Delta_f(a_{2j}, a_{2j-2}). \end{aligned}$$

Observe that $\Delta_f(x, y) + \Delta_f(y, z) = \Delta_f(x, z)$ for all $y \in [x, z]$. Therefore, summation over $j = 1, \dots, 2^{m-1}$ yields

$$(3.3) \quad \sum_{j=1}^{2^m} |f(a_j) - f(a_{j-1})| \leq \sum_{j=1}^{2^{m-1}} |f(a_{2j}) - f(a_{2j-2})| + 2\eta(1) \Delta_f(a, b).$$

Notice that the sum on the right hand side involves only even indices. Next we apply (3.3) to the partition $a_0, a_2, \dots, a_{2^m}$ to rarefy it further. After m steps we arrive at

$$\sum_{j=1}^{2^m} |f(a_j) - f(a_{j-1})| \leq |f(a) - f(b)| + 2m\eta(1) \Delta_f(a, b) \leq (2m+1) |f(a) - f(b)|.$$

This completes the proof of (3.2). $\square$

The following lemma will allow us to derive the conclusion of Theorem 3.1 from the growth estimate (3.2).

Lemma 3.3. *Suppose that $d_1 \geq d_2 \geq \dots \geq d_N$ are positive numbers such that for $j = 1, \dots, N$ the partial sum $s_j := d_1 + \dots + d_j$ is bounded by $C \log(j+1)$, where C is a constant. Let $q > 1$ and the function ϕ be defined by the formula (1.8). Then*

$$(3.4) \quad \sum_{j=1}^N \phi(d_j) \leq C'$$

where C' depends only on C and q .

Proof. Since $d_j \leq C \log(j+1)/j \leq C\sqrt{j}$, it follows that $\log(e+1/d_j) \geq C_1 \log(j+1)$, where C_1 depends only on C . Therefore,

$$(3.5) \quad \sum_{j=1}^N \phi(d_j) \leq C_1^{-q} \sum_{j=1}^N \frac{d_j}{(\log(j+1))^q}.$$

Next we use summation by parts, replacing d_j with $s_j - s_{j-1}$, where $s_0 = 0$ by convention.

$$(3.6) \quad \begin{aligned} & \sum_{j=1}^N \frac{d_j}{(\log(j+1))^q} \\ &= \frac{s_N}{(\log(N+1))^q} + \sum_{j=1}^{N-1} s_j \left(\frac{1}{(\log(j+1))^q} - \frac{1}{(\log(j+2))^q} \right) \end{aligned}$$

The first term on the right is bounded by $C/(\log(N+1))^{q-1}$. Since $s_j \leq C \log(j+1)$ and

$$\frac{1}{(\log(j+1))^q} - \frac{1}{(\log(j+2))^q} \leq \frac{q}{(j+1)(\log(j+1))^{1+q}},$$

it follows that

$$(3.7) \quad \begin{aligned} & \sum_{j=1}^{N-1} s_j \left(\frac{1}{(\log(j+1))^q} - \frac{1}{(\log(j+2))^q} \right) \\ & \leq C \sum_{j=1}^{\infty} \frac{1}{(j+1)(\log(j+1))^q} =: C_2 \end{aligned}$$

where C_2 depends only on C and q . Combining (3.5), (3.6), and (3.7), we obtain (3.4). $\square$

Proof of Theorem 3.1. Let $a, b \in \mathbb{R}^n$ be two distinct points. Given a partition $(a_j)_{j=0}^N$ of $[a, b]$, let $d_1 \geq \dots \geq d_N$ be the numbers $|f(a_j) - f(a_{j-1})|$ arranged in the nonincreasing order. By Lemma 3.2 the partial sums $s_j = d_1 + \dots + d_j$ are bounded by $C \log(2j+2)|f(a) - f(b)|$, where C

is the constant in (3.2). Applying Lemma 3.3 we arrive at the conclusion of the theorem. $\square$

4. FAILURE OF BOUNDED VARIATION: PROOF OF THEOREM 1.5

Proof. Let $Q \subset \mathbb{C}$ be the open square $\{x + iy : 2 < x < 6, |y| < 2\}$. The closure of Q contains two smaller closed squares $Q_1 = \{x + iy : |x - 3| + |y| \leq 1\}$ and $Q_2 = \{x + iy : |x - 5| + |y| \leq 1\}$.




FIGURE 1.

Let $g : \overline{Q} \rightarrow \mathbb{C}$ be a Lipschitz function such that

$$g(z) = \begin{cases} i(z - 2), & z \in Q_1; \\ i(6 - z), & z \in Q_2; \\ 0, & z \in \partial Q. \end{cases}$$

Extend g to the set $A = \bigcup_{k \in \mathbb{Z}} (\overline{Q} + 8k)$ so that $g(z + 8) = g(z)$. Finally, set $g(z) = 0$ for $z \notin A$.

Let L be the Lipschitz constant of g . We shall prove that for $0 < \epsilon < 1/(2L)$ the mapping

$$(4.1) \quad f(z) = z + \epsilon \sum_{m=0}^{\infty} 4^{-m} g(4^m z), \quad z \in \mathbb{C},$$

satisfies

$$(4.2) \quad |f_{\bar{z}}| \leq k \operatorname{Re} f_z \quad \text{a.e.} \quad \text{with} \quad k = \frac{\epsilon L}{1 - \epsilon L}.$$

First of all, the series in (4.1) converges uniformly because g is bounded. Let

$$B = \bigcup_{\ell \in \mathbb{Z}} \{[Q \setminus (Q_1 \cup Q_2)] + 8\ell\}$$

and note that

$$(4.3) \quad B \subset \bigcup_{j \in \mathbb{Z}} \{z: |\operatorname{Re} z - 2j| < |\operatorname{Im} z|\}$$

and

$$(4.4) \quad B \cap \bigcup_{j \in \mathbb{Z}} \{z: |\operatorname{Re} z - 8j| < |\operatorname{Im} z|\} = \emptyset.$$

We claim that for any $z \in \mathbb{C}$ there exists at most one integer $m \geq 0$ such that $4^m z \in B$. Indeed, let m_0 be the smallest such integer. Replacing z with $4^{m_0} z$, we may assume that $m_0 = 0$, i.e., $z \in B$. According to (4.3), there exists $j \in \mathbb{Z}$ such that $|\operatorname{Re} z - 2j| < |\operatorname{Im} z|$. For any $m \geq 1$ the number $\zeta = 4^m z$ satisfies $|\operatorname{Re} \zeta - 8 \cdot 4^{m-1} j| < |\operatorname{Im} \zeta|$, which implies $\zeta \notin B$ by virtue of (4.4). This proves the claim.

Since both $\operatorname{Re} g_z$ and $g_{\bar{z}}$ vanish a.e. outside of B , it follows that

$$1 - \epsilon L \leq \operatorname{Re} f_z(z) \leq 1 + \epsilon L, \quad |f_{\bar{z}}(z)| \leq \epsilon L \quad \text{a.e. in } \mathbb{C}.$$

This proves (4.2). Since $g(z) = 0$ when $|\operatorname{Im} z| > 2$, it follows that $|\operatorname{Im} f_z(z)| \leq \epsilon m L$ when $|\operatorname{Im} z| \geq 2 \cdot 4^{-m}$. This implies

$$(4.5) \quad |Df(z)| \leq C \log(e + 1/|\operatorname{Im} z|)$$

for some constant C . Hence $f \in W_{\text{loc}}^{1,p}(\mathbb{C}; \mathbb{C})$ for all $p < \infty$. Therefore, f is quasiregular. By [13, Cor. 1.5] it is quasiconformal. It is clear that $\operatorname{Re} f(x) = x$ for all $x \in \mathbb{R}$.

It remains to prove that the function

$$h(x) := \epsilon^{-1} \operatorname{Im} f(x) = -i \sum_{m=0}^{\infty} \frac{1}{4^m} g(4^m x), \quad x \in \mathbb{R}$$

has infinite variation on any nontrivial interval $[a, b] \subset \mathbb{R}$. Due to the self-similar structure of h it suffices to consider the interval $[0, 8]$. We will show that the sum

$$V_N := \sum_{j=0}^{4^N} \left| h\left(\frac{8j}{4^N}\right) - h\left(\frac{8j-8}{4^N}\right) \right|$$

satisfies

$$(4.6) \quad V_N \geq c\sqrt{N}$$

with an absolute constant $c > 0$. Let $x_j = (8j)4^{-N}$, $j = 0, \dots, 4^N$. When $m \geq N$, we have $g(4^m x_j) = 0$ for all j . Therefore, in the definition of V_N we can replace h with the partial sum

$$h_N(x) = -i \sum_{m=0}^{N-1} \frac{1}{4^m} g(4^m x).$$

Since h_N is affine on each interval $[x_{j-1}, x_j]$, it follows that $V_N = \int_0^8 |h'_N(x)| dx$.

We claim that for a.e. $x \in \mathbb{R}$

$$(4.7) \quad \frac{d}{dx}(-ig(x)) = \frac{1}{2}(s_0(x/8) - s_1(x/8)),$$

where s_m is the m th Rademacher function, defined by

$$s_m(x) = \text{sign} \sin(2^{m+1}\pi x).$$

Since both sides of (4.7) are periodic functions with period 8, it suffices to check that equality holds a.e. on the interval $(0, 8)$. From the definitions of g and s_m one can see that both sides of (4.7) agree with $\chi_{[2,4]} - \chi_{[4,6]}$ when $0 < x < 8$ and $x \neq 2, 4, 6$. This proves (4.7). It then follows that

$$h'_N(x) = \frac{1}{2} \sum_{m=0}^{N-1} (s_{2m}(x/8) - s_{2m+1}(x/8)).$$

The L^1 norm of a Rademacher series is comparable to the ℓ^2 norm of its coefficients [25, Thm. V.8.4]. Hence $\int_0^8 |h'_N(x)| dx \geq c\sqrt{N}$ with an absolute constant $c > 0$. This completes the proof of (4.6). $\square$

Remark 4.1. The mapping f constructed in the proof of Theorem 1.5 does not have finite ϕ -variation on lines with ϕ as in (1.8) for $0 \leq q < 1/2$.

Proof. Using Jensen's inequality and the estimate (4.6), we obtain

$$\begin{aligned} \sum_{j=0}^{4^N} \phi(|h(x_j) - h(x_{j-1})|) &\geq 4^N \phi(V_N/4^N) \\ &\geq \frac{c\sqrt{N}}{(\log(e + 4^N/(c\sqrt{N})))^q} \rightarrow \infty \end{aligned}$$

as $N \rightarrow \infty$. $\square$

Proof of Corollary 1.6. We may assume that the lines L_j are parallel to the real axis; that is, $L_j = \{z \in \mathbb{C} : \text{Im } z = b_j\}$ where b_j are distinct real numbers. For $m = 1, 2, \dots$ let $\epsilon_m = \min_{1 \leq j < \ell \leq m} |b_j - b_\ell|$ and choose $c_m > 0$ so that

$$(4.8) \quad c_m |b_m| < 2^{-m}$$

and

$$(4.9) \quad c_m \log(e + 1/\epsilon_m) < 2^{-m}.$$

We define

$$(4.10) \quad F(z) = \sum_{m=1}^{\infty} c_m f(z - ib_m)$$

where f is the mapping in (4.1). Note that $|f(z)| \leq |z| + M$ for some constant M . The sum in (4.10) converges locally uniformly because by (4.8)–(4.9)

$$|c_m f(z - ib_m)| \leq c_m (|z| + |b_m| + M) \leq 2^{-m} (|z| + 1 + M).$$

Therefore F is quasiconformal [15, II 5.3]. We claim for every $j = 1, 2, \dots$ the sum

$$R_j(z) = \sum_{m \neq j} c_m f(z - ib_m)$$

is Lipschitz on the line L_j . By (4.5) the restriction of $f(z - ib_m)$ to L_j is Lipschitz with a constant $C \log(e + 1/|b_m - b_j|)$. We estimate the Lipschitz constant of R_j by

$$\sum_{m \neq j} c_m \log(e + 1/|b_m - b_j|) \leq \sum_{m < j} c_m \log(e + 1/\epsilon_j) + \sum_{m > j} c_m \log(e + 1/\epsilon_m).$$

The first sum on the right has finitely many terms, and the second sum converges by (4.9). Since $F(z) = c_j f(z - ib_j) + R_j$, it follows that $F(L_j)$ is not locally rectifiable at any of its points. $\square$

5. REDUCED QUASICONFORMAL MAPPINGS IN FOUR DIMENSIONS

Our first goal in this section is to reformulate the definition of reduced quasiconformal mappings (Definition 1.4) in terms of differential matrices $Df(x) \in \mathbb{M}_n$ rather than complex derivatives. This is done in Proposition 5.3 below. We write $\mathbb{M}_n$ for the set of all $n \times n$ matrices with real entries. Also, for $\delta \in [-1, 1]$ we let

$$\mathbb{M}_n(\delta) = \{A \in \mathbb{M}_n : \langle Av, v \rangle \geq \delta |Av| |v| \text{ for all } v \in \mathbb{R}^n\}.$$

Proposition 5.1. *For $\delta \in (0, 1)$ let $H(\delta) = \frac{1+\sqrt{1-\delta^2}}{1-\sqrt{1-\delta^2}}$. Then $\|A\| \|A^{-1}\| \leq H(\delta)$ for all nonzero matrices $A \in \mathbb{M}_n(\delta)$.*

Proof. For $n = 2$ this proposition was proved in [2, p.84]. If $n \geq 3$, let v and w be distinct unit vectors such that $|Av| = \|A\|$ and $|Aw| = \|A^{-1}\|^{-1}$. Applying the two-dimensional case to the subspace spanned by v and w , we arrive at the desired conclusion. $\square$

A matrix $A \in \mathbb{M}_2$ determines a linear mapping $x \mapsto Ax$ of the plane $\mathbb{R}^2$. The same linear mapping can be written as $z \mapsto \alpha^+ z + \alpha^- \bar{z}$ for some $\alpha^+, \alpha^- \in \mathbb{C}$. The numbers α^+ and α^- can be thought of as the conformal and anticonformal parts of A (cf. [8]).

Proposition 5.2. [2, Thm. 3.11.6] *A matrix A belongs to $\mathbb{M}_2(\delta)$ if and only if $|\alpha^-| + \delta |\operatorname{Im} \alpha^+| \leq \sqrt{1 - \delta^2} \operatorname{Re} \alpha^+$.*

A complex number $a + ib \in \mathbb{C}$ can be identified with the 2×2 matrix $Z = \begin{pmatrix} a & -b \\ b & a \end{pmatrix}$. Thus we may consider $\mathbb{C}$ as a linear subspace of $\mathbb{M}_2$. Within this subspace each matrix decomposes into real and imaginary parts: $\operatorname{Re} Z = \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}$ and $\operatorname{Im} Z = \begin{pmatrix} 0 & -b \\ b & 0 \end{pmatrix}$. Given $A \in \mathbb{M}_2$, let $\mathbb{C}(A)$ be the orthogonal projection of A onto the subspace $\mathbb{C}$.

Proposition 5.3. *A mapping $f \in W_{\text{loc}}^{1,2}(\mathbb{R}^2; \mathbb{R}^2)$ is reduced quasiconformal in the sense of Definition 1.4 if and only if there exists $\delta > 0$ such that for a.e. $x \in \mathbb{R}^2$ the derivative $A = Df(x)$ satisfies $A - \operatorname{Im} \mathbb{C}(A) \in \mathbb{M}_2(\delta)$.*

Proof. Writing the matrix $A = Df(x)$ in conformal-anticonformal coordinates as (α^+, α^-) , we observe that $A - \operatorname{Im} \mathbb{C}(A)$ corresponds to $(\operatorname{Re} \alpha^+, \alpha^-)$. According to Proposition 5.2, the condition $A - \operatorname{Im} \mathbb{C}(A) \in \mathbb{M}_2(\delta)$ is equivalent to $|\alpha^-| \leq \sqrt{1 - \delta^2} \operatorname{Re} \alpha^+$. The latter inequality is the same as (1.6) with $k = \sqrt{1 - \delta^2}$. $\square$

A quaternion $\alpha + \beta \mathbf{i} + \gamma \mathbf{j} + \zeta \mathbf{k}$ can be identified with a 4×4 real matrix

$$(5.1) \quad Q = \begin{pmatrix} \alpha & -\beta & -\gamma & -\zeta \\ \beta & \alpha & -\zeta & \gamma \\ \gamma & \zeta & \alpha & -\beta \\ \zeta & -\gamma & \beta & \alpha \end{pmatrix}$$

With this identification we consider the set of quaternions $\mathbb{H}$ as a subset of $\mathbb{M}_4$. Since quaternion conjugation corresponds to matrix transposition, we have $Q^T Q = \|Q\|^2 I$, where $\|Q\|$ is the operator norm of matrix Q , also equal to the absolute value of the quaternion. Consequently, $|Qv| = \|Q\| |v|$ for any vector $v \in \mathbb{R}^4$.

A quaternion Q is the sum of its real (scalar) and imaginary parts:

$$(5.2) \quad \operatorname{Re} Q = \begin{pmatrix} \alpha & 0 & 0 & 0 \\ 0 & \alpha & 0 & 0 \\ 0 & 0 & \alpha & 0 \\ 0 & 0 & 0 & \alpha \end{pmatrix}, \quad \operatorname{Im} Q = Q - \operatorname{Re} Q.$$

If $\operatorname{Re} Q = 0$, the quaternion Q is purely imaginary. For a matrix $A \in \mathbb{M}_4$, we define $\mathbb{H}(A)$ to be the orthogonal projection of A onto the subspace $\mathbb{H} \subset \mathbb{M}_4$.

Definition 5.4. A homeomorphic mapping $f \in W_{\text{loc}}^{1,4}(\mathbb{R}^4; \mathbb{R}^4)$ is *reduced quasiconformal* if there exists $\delta > 0$ such that for a.e. $x \in \mathbb{R}^4$ the derivative $A = Df(x)$ satisfies $A - \operatorname{Im} \mathbb{H}(A) \in \mathbb{M}_4(\delta)$.

First of all, we need to justify the terminology by proving the following proposition.

Proposition 5.5. *Any reduced quasiconformal mapping $f: \mathbb{R}^4 \rightarrow \mathbb{R}^4$ is K -quasiconformal, where K depends only on δ in Definition 5.4. In addition, f is monotone.*

Proof. The essence of this proposition is the algebraic implication

$$(5.3) \quad A - \operatorname{Im} \mathbb{H}(A) \in \mathbb{M}_4(\delta) \implies \|A\| \|A^{-1}\| \leq \tilde{H}(\delta).$$

We may assume that A is a nonzero matrix. Let $Q = \operatorname{Im} \mathbb{H}(A)$ and $B = A - Q$. If $Q = 0$, then Proposition 5.1 gives (5.3) with $\tilde{H}(\delta) = H(\delta)$. Assume $Q \neq 0$. Fix a unit vector $v \in \mathbb{R}^4$. Since $Qv/\|Q\|$ is a unit vector orthogonal to v , it follows that

$$\langle Bv, v \rangle^2 + \|Q\|^{-2} \langle Bv, Qv \rangle^2 \leq |Bv|^2$$

Using the inequality $\langle Bv, v \rangle \geq \delta |Bv|$, we obtain

$$(5.4) \quad |\langle Bv, Qv \rangle| \leq \sqrt{1 - \delta^2} \|Q\| |Bv|,$$

which in turn yields

$$(5.5) \quad \begin{aligned} |Av|^2 &= |Bv + Qv|^2 = |Bv|^2 + \|Q\|^2 + 2\langle Bv, Qv \rangle \\ &\geq (1 - \sqrt{1 - \delta^2})(|Bv|^2 + \|Q\|^2) + \sqrt{1 - \delta^2}(|Bv| - \|Q\|)^2 \\ &\geq (1 - \sqrt{1 - \delta^2})(|Bv|^2 + \|Q\|^2). \end{aligned}$$

In particular, A is invertible. We also have the trivial estimate

$$(5.6) \quad |Av|^2 \leq 2(\|Bv\|^2 + \|Q\|^2).$$

Combining (5.5) and (5.6), we conclude that

$$\begin{aligned} \|A\|^2 \|A^{-1}\|^2 &= \frac{\max\{|Av|^2 : |v| = 1\}}{\min\{|Av|^2 : |v| = 1\}} \\ &\leq \frac{2}{1 - \sqrt{1 - \delta^2}} \frac{\max\{|Bv|^2 : |v| = 1\} + \|Q\|^2}{\min\{|Bv|^2 : |v| = 1\} + \|Q\|^2} \\ &\leq \frac{2}{1 - \sqrt{1 - \delta^2}} \frac{\max\{|Bv|^2 : |v| = 1\}}{\min\{|Bv|^2 : |v| = 1\}} \leq \frac{2(1 + \sqrt{1 - \delta^2})^2}{(1 - \sqrt{1 - \delta^2})^3} \end{aligned}$$

where the last step uses Proposition 5.1. This proves (5.3). Applying (5.3) to the derivative matrix $A = Df(x)$, we find that f is K -quasiconformal with $K = \tilde{H}(\delta)^3$.

With $A = Df(x)$ and $B = A - \operatorname{Im} \mathbb{H}(A)$ as above, we have

$$\langle Av, v \rangle = \langle Bv, v \rangle \geq 0, \quad v \in \mathbb{R}^4.$$

Integrating this inequality along the segments $[a, b]$ on which f is absolutely continuous, we obtain $\Delta_f(a, b) \geq 0$. The continuity of f then implies $\Delta_f(a, b) \geq 0$ for all $a, b \in \mathbb{R}^4$. $\square$

Remark 5.6. If in Definition 5.4 we do not require f to be homeomorphic, then the proof of Proposition 5.5 shows that f is K -quasiregular (see Definition 1.2).

It follows from Definition 5.4 that the set of reduced quasiconformal mappings is a convex cone in four dimensions as well as in two dimensions. Another similarity with the planar case is provided by the following result.

Proposition 5.7. *Any nonconstant δ -monotone mapping $f: \mathbb{R}^4 \rightarrow \mathbb{R}^4$ is reduced quasiconformal in the sense of Definition 5.4.*

Proof. Since f is quasiconformal by [16, Cor. 7], we have $f \in W_{\text{loc}}^{1,4}(\Omega; \mathbb{R}^4)$. Fix a point $x \in \mathbb{R}^4$ where f is differentiable and let $A = Df(x)$, $Q = \text{Im } \mathbb{H}(A)$, $B = A - Q$. The definition of δ -monotonicity implies $A \in \mathbb{M}_4(\delta)$. For all $v \in \mathbb{R}^n$ we have $\langle Qv, v \rangle = 0$ since Q is an antisymmetric matrix. Thus,

$$(5.7) \quad \langle Bv, v \rangle = \langle Av, v \rangle \geq \delta |Av| |v|, \quad v \in \mathbb{R}^n.$$

It remains to prove that $|Av| \geq c |Bv|$ for a constant $c > 0$ that depends only on δ . Note that $\text{Im } \mathbb{H}(A)$ is the orthogonal projection of A onto the space of purely imaginary quaternions, considered as a linear subspace of $\mathbb{M}_4$. Therefore, $B = A - \text{Im } \mathbb{H}(A)$ is the projection of A onto the orthogonal complement of purely imaginary quaternions. Since orthogonal projections in $\mathbb{M}_n$ do not increase the Frobenius norm $\|\cdot\|_F$, it follows that

$$(5.8) \quad \|B\| \leq \|B\|_F \leq \|A\|_F \leq 2\|A\|$$

where we have used the relation between operator norm and Frobenius norm [12, p.313]. Combining Proposition 5.1 with (5.8) we obtain

$$|Av| \geq \frac{\|A\| |v|}{H(\delta)} \geq \frac{\|B\| |v|}{2H(\delta)} \geq \frac{|Bv|}{2H(\delta)}.$$

This estimate together with (5.7) imply $B \in \mathbb{M}_4(\delta/2H(\delta))$. $\square$

Our last result is an extension of Theorem 1.7 to four dimensions.

Theorem 5.8. *Let $f: \mathbb{R}^4 \rightarrow \mathbb{R}^4$ be a reduced quasiconformal mapping in the sense of Definition 5.4. Then for any $q > 1$ f has finite ϕ -variation on lines with ϕ as in (1.8).*

Proof. For a purely imaginary quaternion $Q \in \mathbb{M}_4$ we define $f^Q(x) = f(x) + Qx$. Recall the definition of the modulus of monotonicity Δ_f in (2.3). We claim that

$$(5.9) \quad \Delta_f(a, b) = \min_Q |f^Q(a) - f^Q(b)|$$

where the minimum is taken over all purely imaginary quaternions (and is attained). Indeed,

$$\Delta_f(a, b) = \Delta_{f^Q}(a, b) \leq |f^Q(a) - f^Q(b)|, \quad \text{for any } Q \text{ with } \text{Re } Q = 0.$$

In proving the converse inequality we may assume that $v := a - b$ is a nonzero vector. Applying the unit quaternions $\mathbf{i}$, $\mathbf{j}$, and $\mathbf{k}$ to v , we obtain an

orthogonal basis of $\mathbb{R}^4$, namely $\{v, \mathbf{i}v, \mathbf{j}v, \mathbf{k}v\}$. Expand the vector $f(a) - f(b)$ in this basis:

$$(5.10) \quad f(a) - f(b) = \alpha v + \beta \mathbf{i}v + \gamma \mathbf{j}v + \zeta \mathbf{k}v.$$

In these terms, $\Delta_f(a, b) = \langle \alpha v, v/|v| \rangle = \alpha|v|$. Since f is monotone by Proposition 5.5, we have $\alpha \geq 0$. The quaternion $Q = -\beta \mathbf{i} - \gamma \mathbf{j} - \zeta \mathbf{k}$ satisfies

$$|f^Q(a) - f^Q(b)| = \alpha|v| = \Delta_f(a, b)$$

which proves (5.9). For future reference, observe that (5.10) implies $|f(a) - f(b)| \geq |Qv| = \|Q\||v|$, hence

$$(5.11) \quad \|Q\| \leq \frac{|f(a) - f(b)|}{|a - b|}.$$

Since linear mappings trivially satisfy the conclusion of the theorem, we may assume that f is nonlinear. By Remark 5.6 there exist $K < \infty$ such that f^Q is K -quasiregular for all purely imaginary quaternions. Also, f^Q is monotone by Proposition 5.5. By Theorem 1.2 [17] any monotone quasiregular mapping defined on $\mathbb{R}^n$ is either constant or a homeomorphism. Since f is not linear, f^Q cannot be constant. Thus f^Q is K -quasiconformal. By [11, Thm 11.14] the family f^Q has a common modulus of quasismmetry η . Given distinct points $a, b, c \in \mathbb{R}^4$, let Q be a minimizing quaternion in (5.9). The quasismmetry of f^Q implies

$$|f^Q(c) - f^Q(a)| \leq \eta \left(\frac{|c - a|}{|b - a|} \right) |f^Q(b) - f^Q(a)|.$$

Here $|f^Q(b) - f^Q(a)| = \Delta_f(a, b)$. Using (5.11) we obtain

$$\begin{aligned} |f(c) - f(a)| &\leq |Q(c - a)| + |f^Q(c) - f^Q(a)| \\ &\leq \frac{|c - a|}{|a - b|} |f(a) - f(b)| + \eta \left(\frac{|c - a|}{|b - a|} \right) \Delta_f(a, b), \end{aligned}$$

which is (3.1). It remains to apply Theorem 3.1 to f . $\square$

ACKNOWLEDGEMENT

We thank Tadeusz Iwaniec for many stimulating discussions on the subject of this paper.

REFERENCES

1. G. Alessandrini and V. Nesi, *Beltrami operators, non-symmetric elliptic equations and quantitative Jacobian bounds*, Ann. Acad. Sci. Fenn. Math., to appear.
2. K. Astala, T. Iwaniec, and G. J. Martin, *Elliptic partial differential equations and quasiconformal mappings in the plane*, Princeton Mathematical Series, vol. 48, Princeton University Press, 2009.
3. K. Astala and J. Jääskeläinen, *Homeomorphic solutions to reduced Beltrami equations*, preprint, arXiv:0812.2322.

4. Z. Balogh, P. Koskela, and S. Rogovin, *Absolute continuity of quasiconformal mappings on curves*, Geom. Funct. Anal. **17** (2007), no. 3, 645–664.
5. C. J. Bishop, *Quasiconformal mappings which increase dimension*, Ann. Acad. Sci. Fenn. Math. **24** (1999), no. 2, 397–407.
6. M. Bonk, J. Heinonen, and E. Saksman, *The quasiconformal Jacobian problem*, in “In the tradition of Ahlfors and Bers. III”, 77–96, Contemp. Math., 355, Amer. Math. Soc., Providence, RI, 2004.
7. L. C. Evans and R. F. Gariepy, *Measure theory and fine properties of functions*, CRC Press, Boca Raton, FL, 1992.
8. D. Faraco and L. Székelyhidi, *Tartar’s conjecture and localization of the quasiconvex hull in $\mathbb{R}^{2 \times 2}$* , Acta Math. **200** (2008), no. 2, 279–305.
9. F. W. Gehring, *The definitions and exceptional sets for quasiconformal mappings*, Ann. Acad. Sci. Fenn. Ser. AI Math. **281** (1960), 1–28.
10. F. W. Gehring, *The L^p -integrability of the partial derivatives of a quasiconformal mapping*, Acta Math. **130** (1973), 265–277.
11. J. Heinonen, *Lectures on analysis on metric spaces*, Universitext. Springer-Verlag, New York, 2001.
12. R. A. Horn and C. R. Johnson, *Matrix analysis*, Cambridge University Press, Cambridge, 1985.
13. T. Iwaniec, L. V. Kovalev, and J. Onninen, *On injectivity of quasiregular mappings*, Proc. Amer. Math. Soc., to appear.
14. T. Iwaniec, L. V. Kovalev, and J. Onninen, *Dynamics of quasiconformal fields*, preprint, arXiv:0811.4217.
15. O. Lehto and K. I. Virtanen, *Quasiconformal mappings in the plane*, 2nd edition. Springer-Verlag, New York–Heidelberg, 1973.
16. L. V. Kovalev, *Quasiconformal geometry of monotone mappings*, J. Lond. Math. Soc. (2) **75** (2007), no. 2, 391–408.
17. L. V. Kovalev and J. Onninen, *On invertibility of Sobolev mappings*, preprint, arXiv:0812.2350.
18. J. Malý, *Absolutely continuous functions of several variables*, J. Math. Anal. Appl. **231** (1999), no. 2, 492–508.
19. J. Musielak and W. Orlicz, *On generalized variations. I*, Studia Math. **18** (1959) 11–41.
20. Yu. G. Reshetnyak, *Space mappings with bounded distortion*, American Mathematical Society, Providence, RI, 1989.
21. S. Rickman, *Quasiregular mappings*, Springer-Verlag, Berlin, 1993.
22. P. Tukia, *A quasiconformal group not isomorphic to a Möbius group*, Ann. Acad. Sci. Fenn. Ser. A I Math. **6** (1981), no. 1, 149–160.
23. D. Waterman, *On convergence of Fourier series of functions of generalized bounded variation*, Studia Math. **44** (1972), 107–117.
24. L. C. Young, *General inequalities for Stieltjes integrals and the convergence of Fourier series*, Math. Ann. **115** (1938), no. 1, 581–612.
25. A. Zygmund, *Trigonometric series*, 3rd edition. Cambridge University Press, Cambridge, 2002.

DEPARTMENT OF MATHEMATICS, SYRACUSE UNIVERSITY, 215 CARNEGIE, SYRACUSE, NY 13244, USA

E-mail address: `lvkovale@syr.edu`

DEPARTMENT OF MATHEMATICS, SYRACUSE UNIVERSITY, 215 CARNEGIE, SYRACUSE, NY 13244, USA

E-mail address: `jkonnine@syr.edu`